

(RESEARCH ARTICLE)



## Trace metals assessment in agricultural soil and irrigation water at hail region Saudi Arabia

Amal H Al-Bagawi \*, Dorsaf Mansour and Shomokh AM Aljabri

*Department of Chemistry, Faculty of Science, University of Ha'il, Hail 1560, KSA.*

International Journal of Frontline Research in Chemistry and Pharmacy, 2022, 01(01), 021–031

Publication history: Received on 26 February 2022; revised on 02 April 2022; accepted on 04 April 2022

Article DOI: <https://doi.org/10.56355/ijfrcp.2022.1.1.0004>

### Abstract

The suitability of agricultural soil and irrigation water depends on the presence of trace metals; the aim of the present work is to assessment of the trace metals at Hail region using analytical techniques. The metal contaminations levels were determined using ICP-MS and flame photometer. The result obtained were compared to the standard samples suitable for agricultural and it was in the suitable range. The geo-accumulation (I-geo), single pollution, Nemerow pollution indices showed that the Se pollution intensity was significant for agricultural soils. I-geo values revealed no real sign of contamination with almost all the samples, reflecting a lack of contamination for all elements except Zn. While the enrichment factor (EF) for Fe was less than 2 suggested that the elements come entirely from crustal materials or natural processes. The study showed considerable variation in the levels of the analyzed elements in the soil samples. The total metal concentrations in the soil samples ordered as follows: Na > K > Fe > Mn > Zn > Ba > Cr > Ni > Cu > Pb > Cd > As.

**Keywords:** Contamination assessment; Water; Agricultural soil; ICP-MS; Trace metals; Saudi Arabia; Hail region

### 1. Introduction

The suitability of water and soil for agriculture depend on the presence and concentrations of some elements, specially the heavy metals. In the last years and due to increment of the human activities the water pollution by heavy metals increased rapidly and the estimation of this metals periodically is so important process to understand the suitability of the irrigation water and soil for the agriculture. Heavy metals are widespread pollutants of great environmental concern as they are non degradable, toxic, and persistent [1-3]. Heavy metals and some trace elements are biologically toxic and can affect the health of human being owing to their accumulation and persistence in the compartments of the food chain. Heavy metals become toxic at a certain concentrations and when they are not metabolized in the body and accumulate in the soft tissues [4-5]. The area of the research in the a quatic chemistry of some trace elements is sovibrant areas aand considerd as a very needing research area with a wide spectrum of aspects and considrations to be studied for the soil and water. The determination of trace elements in water and soil is of widespread and great importance. Trace elements can be categorized as (1) essential to human life (chromium, copper, iodine, molybdenum, selenium, and zinc), (2, 3) probably essential (boron, manganese, nickel, silicon, and vanadium), and potentially toxic, some of which possibly have essential functions (aluminum, arsenic, cadmium, fluoride, lead, lithium, mercury, and tin) [3]. Groundwater is one of the most important natural resources for domestic, agriculture and industrial uses in an arid country like Saudi Arabia.

The previous study on the Soil and water samples at Hail Region KSA, was conducted at 2016;s from about 5 years ago and not covered all Hail areas. During that period of time the concentrations of trace elements have been changed so the periodic estimation of the trace metals is tremndous .

\* Corresponding author: Amal H Al-Bagawi  
Department of Chemistry, Faculty of Science, University of Ha'il, Hail 1560, KSA.

Analytical and electroanalytical techniques are highly effective for the trace elements determination [13-22]. From the most popular analytical techniques are Atomic absorption spectroscopy, AAS, gravimetric, complexometric, ICP-MS, and UV- Spectroscopy [13-22]. The most popular electrochemical method for determination of trace elements in water samples, are the potentiometry and voltammetry. This method is very sensitive for many elemental analysis, and can be used for simultaneous determination at low cost. In particular, it is favorable for the analysis of many elements like cadmium, copper, lead and zinc [7-9]. So, the aim of the present research is to investigate the distribution of some elements in the irrigation water and agricultural soil at Hail region to assess the element contamination levels and to reach the suitability of the water and soil for agriculture by through comparison models studies and using analytical techniques. The analytical techniques which will be used in this study are Atomic absorption spectroscopy, AAS, and ICP-MS. The results were compared according to soil quality guidelines. Both geo-accumulation (I-geo), single pollution, and Nemerow pollution will be used to gain information about the sources of metal pollutants and to assess the metal pollution status. All the data obtained from the used different techniques were compared and discussed in order to determine the suitability of the water and soil for agriculture at Hail Region KSA.

## 2. Material and methods

### 2.1 The Area of the Study

This study was conducted during 2020 in agricultural areas at Hail region, and its Villages [23]. The average annual rainfall in study location is about 10.16 cm without any, or with a negligible amount, of rains in the summer season [24].

### 2.2 Water sample collection and analysis

Ground water samples used for irrigation were collected and preserved according to the operating procedure for groundwater sampling [23-25]. A total of fifty samples were brought from several farms within the Hail area. Well Tight-capped high quality polyethylene terephthalate bottles were used for sample storage. Before use, the bottles were washed by distilled deionized water and rinsed overnight in 10% (v/v) nitric acid. Samples were filtered through the Whatmann filter papers number 42. To prevent precipitation of metals and biological growth, few drops of concentrated nitric acid were added to samples to obtain pH nearby 2 [26,27]. Thereafter, samples were immediately transported to the laboratory in iceboxes and stored at 4 °C up to analysis. The elements were measured using Thermo Scientific iCAPQ ICP-MS with CETAC ASX-520 Autosampler). The precision of the water method analysis for the multi-element determination was evaluated by using triplicate readings for each analysis, and the mean was calculated with relative standard deviations less than 4%.

### 2.3 Soil samples collection and analysis

A total of 50 soil samples were collected from agricultural farms situated along the main highways in the Hail region and her Villages during 2018. The samples were collected from the upper 10 cm sections. After grinding and sieving through 2 mm mesh, 500 mg were digested according to [26, 27]. The elements were measured using an ICP-MS: With Auto Sampler ICAP Q, CETAC ASX-520, Thermo scientific USA. The precision of the soil method analysis for the multi-element determination was evaluated by using triplicate readings for each analysis, and the mean was calculated with relative standard deviations less than 4%.

### 2.4 Contamination level of elements

I-geo, pollution load index (PLI), and enrichment factors were used to gain information about the sources of metal pollutants and to assess the metal pollution status.

#### 2.4.1 Pollution Load Index

PLI was evaluated as follows equation (1) [25-28];

$$PLI = (P_{i1} \times P_{i2} \times P_{i3} \times \dots \times P_{in})^{1/n} \quad (1)$$

Where n is the number of elements, and  $P_i$  is the single pollution index by element i, it is the ratio between the element level ( $C_i$ ) in soil samples and its background concentration ( $S_i$ ) and given by the following equation (2):

$$P_i = C_i/S_i \quad (2)$$

Where PLI value >1 would indicate a contaminated site while PLI value <1 indicates no contamination.

#### 2.4.2 Geo-accumulation index

*I-geo* was computed as follows, equation (3) [28,29].

$$I\text{-geo} = \text{Log}_2 (C_n / 1.5B_n) \quad (3)$$

Where  $C_n$  is the total metal concentration in the soil sample;  $B_n$  is the metal background value, and the value 1.5 represent the factor for background matrix correction. *I-geo* consists of seven classes as shown in Table 1. The concentrations ( $\mu\text{g/g}$ ) of Zn (200), Cu (63), Pb (70), Ni (45), Co (40), Cr (64), Se (1) and Cd (1.4) in the Canadian soil quality guidelines [21], were used as background.

**Table 1** Descriptive classes for  $I_{\text{geo}}$  values

$I_{\text{geo}}$ Class	0	1	2	3	4	5	6
$I_{\text{geo}}$	$I_{\text{geo}} < 0$	$0 < I_{\text{geo}} \leq 1$	$1 < I_{\text{geo}} \leq 2$	$2 < I_{\text{geo}} \leq 3$	$3 < I_{\text{geo}} \leq 4$	$4 < I_{\text{geo}} \leq 5$	$I_{\text{geo}} > 5$
Sediment quality	Unpolluted	Unpolluted to moderately polluted	Moderately polluted	Moderately to heavily polluted	Heavily polluted	Heavily to extremely polluted	Extremely polluted

#### 2.4.3 Nemerow pollution index

The Nemerow integrated pollution index  $P$  considers the average values of all studied elements in addition to the maximum value of  $P_i$ , where  $P_{\text{ave}}$  and  $P_{\text{max}}$  represent the average and maximum of the pollution indices for each element, respectively [29]. The following equation determines  $P$ : Equation (3) [22].

$$P = \sqrt{\frac{P_{\text{max}}^2 + P_{\text{ave}}^2}{2}} \quad (4)$$

#### 2.4.4 Enrichment Factor

The EF of a single trace element in the soils was calculated as follows equation (5) [30].

$$EF = \frac{\left(\frac{M}{Al}\right)_{\text{sample}}}{\left(\frac{M}{Al}\right)_{\text{background}}} \quad (5)$$

Al was used as a conservative tracer to differentiate natural from anthropogenic components.  $(M/Al)_{\text{sample}}$  is the ratio of metal, and Al concentration in the sample in the examined environment and  $(M/Al)_{\text{background}}$  is the ratio of metal and Al concentration of the background [24]. The background concentrations of Fe, Mn, Zn, Cu Ni, Cr, Pb, Cd, Co and Se in the Upper Continental Crust obtained from TAYLOR and MCLENNAN [31] were used. EF values were classified as  $EF < 2$ , clean-light pollution;  $2 \leq EF < 5$ , moderate pollution;  $5 \leq EF < 20$ , significant pollution;  $20 \leq EF < 40$ , strong pollution;  $EF \geq 40$ , extreme pollution. Data of water and soil samples were examined for significant differences for all studied elements fractions among different locations by ANOVA test.

## 3. Results and discussion

### 3.1 Metals Concentrations in Hail irrigation water

The concentrations of trace elements in the hail irrigation water was measured by ICP-MS analytical techniques for all elements except Na and K which were measured using the flame photometer all concentrations are mentioned in  $\mu\text{g/L}$ . the data obtained were summarized in table 2. As show in table 2 that: the concentrations ( $\mu\text{g/L}$ ) of elements in the ground water used for irrigation in Hail area. No significant differences ( $P > 0.05$ ) in the water trace elements concentration were observed between the studied sites along Hail region. The perusal water data revealed that, the trend of elements according to mean concentration in the samples was:  $\text{Na} > \text{K} > \text{Mn} > \text{Ba} > \text{Fe} > \text{Zn} > \text{Pb} > \text{Cr} > \text{Ni} > \text{Cu} > \text{Cd} > \text{As}$ . The variation in trace elements concentration is controlled by the variation in local and regional geology, water/rock interactions, dilution due to precipitation [2].

**Table 2** Summary of elements concentrations  $\mu\text{g/L}$  in Hail irrigation water

Variable	Max	Min	Median	Mean	SD	CCME (2007)
Na	3670	16	418	819.0571	1055.701	-
K	68.1	1.4	7.5	11.7532	15.61767	-
As	0.001	0.001	0.001	0.001	$4.44 \times 10^{-19}$	100
Cd	0.001	0.001	0.001	0.001	$4.44 \times 10^{-19}$	5.1
Pb	0.036	0.001	0.001	0.001	0.007638	200
Cr	0.0138	0.001	0.001	0.001	0.004258	8
Ni	0.001	0.001	0.001	0.001	$4.44 \times 10^{-19}$	200
Fe	0.07941	0.0057	0.00314	0.00997	0.018668	5000
Mn	13.164	0.00501	0.001	0.628115	2.872332	200
Ba	0.0819	0.00504	0.0319	0.034799	0.019752	100
Cu	0.001	0.001	0.001	0.001	$4.44 \times 10^{-19}$	-
Zn	0.04415	0.0058	0.001	0.005285	0.010925	-

### 3.2 Metals Concentrations in Hail soil

The concentrations of the trace metals in the agricultural soil were measured by ICP-MS analytical techniques for all elements except Na and K which were measured using the flame photometer all concentrations are mentioned in  $\mu\text{g/L}$ . The data obtained were summarized in table 3. As shown in table 3 that: the trend of elements according to mean concentration was:  $\text{Na} > \text{K} > \text{Fe} > \text{Mn} > \text{Zn} > \text{Ba} > \text{Cr} > \text{Ni} > \text{Cu} > \text{Pb} > \text{Cd} > \text{As}$ . The variation in trace elements concentration in Hail region may be due to irrigation of land by fertilizers and other agronomic practices containing metals. Across the investigated samples, wide elements concentration ranges have been recorded in Table 3. No significant differences ( $P > 0.05$ ) in the soil trace elements concentration were observed between the studied sites along Hail region. Soil contamination with toxic and trace elements due to point sources or parent materials often occurs and is easy to identify. The use of elements-enriched chemicals, fertilizers, and organic amendments such as sewage sludge as well as wastewater may cause contamination at a large scale [32]. The higher concentration of Na and K is normal which result due to the nature of soil and the decay of the plants tissue, papers, and routes. Fe is the fourth most abundant element in the Earth's crust. It is the most predominant among studied elements in the Hail region and varied in the range of 8.37–29.89 mg/g. The Hail soil had distinctive red color owing to the occurrence of iron oxides [33], so the Fe concentrations of a soil are region specific and can vary considerably locally due to soil types. Various results revealed the very high levels of Fe in soils and concluded their carcinogenic/mutagenic effects on the living being's health [34]. Fe and Mn occur naturally at plentiful levels, thus are rarely affected by anthropogenic inputs. For agricultural considerations, higher tissue levels of manganese are usually found in the older leaves on the plant and may be associated with damaged or diseased leaves. Mn varied in the range of 0.10–4.77 mg/g. It is considered an essential metal to controlling the behavior of several micronutrients in the soil, but it may cause severe problems if found in high levels. Various reports indicating the high application of Mn in agriculture are available [35]. Although Zn is an essential trace element, high levels can cause harmful health effects. Zn varied in the range of 43.29–1293  $\mu\text{g/g}$ , where 70% of samples were estimated to be more than the maximum permissible level cited by CCME, indicating that there is relatively Zn pollution and relating to the application of Zn fertilizer. Barium (Ba) concentrations range was 21.21  $\mu\text{g/L}$  in the irrigation water samples. The Ba concentrations higher than the CCME limit (100  $\mu\text{g/L}$ ) for irrigation water. The US Environmental Protection Agency found that acute exposure to Ba at above the maximum contaminant level (2 mg/L) can potentially cause gastrointestinal disturbances and muscular weakness. Long-term exposure to Ba at can also potentially cause hypertension. There is no evidence that lifetime exposure to Ba in drinking water has the potential to cause cancer [33]. Other elements from Cr to As ( $\text{Cr} > \text{Ni} > \text{Cu} > \text{Pb} > \text{Cd} > \text{As}$ ) were exist in the trace concentrations. The mean values of Cr, Ni, Cu, Pb, Cd, and As did not exceed the Canadian soil guidelines, which mean that the soil is not contaminated. Zn were present in high levels and had a significant contamination level compared to other elements.

**Table 3** Summary of elements concentration ( $\mu\text{g/g}$ ) in Hail soil

Analytical techniques	Data obtained from Flame photometer		Data obtained from ICP-MS									
	Na	K	As	Cd	Pb	Cr	Ni	Fe	Mn	Ba	Cu	Zn
Max	1869	825	0.001	0.2611	5.152	15.715	14.2205	5145	137.865	21.21	7.588	75.495
Min	24	54	0.001	0.001	0.31591	1.93445	1.12665	738.85	10.752	2.6124	0.57645	1.0913
Median	135	297	0.001	0.00791	1.26525	6.1495	5.4775	2906.4	84.77	13.2685	2.76675	28.882
Mean	241.133	360	0.001	0.02721	1.57276	6.5997	5.76274	2825.75	73.1469	12.1277	3.18398	29.1399
SD	532.734	237.583		0.0241	0.76183	3.28753	3.0612	711.13	29.5262	4.89786	1.43869	16.6138
CCME	45	-	100	104	70	64	45	5000	200	100	63	200

### 3.3 Comparing the trace elements in the study area to the soil in other regions in Saudi Arabia

The present results recorded high levels of Zn, Cu, Ni, and Cr than the other regions (Table 4). The abundance of individual elements in soils and other surficial materials e.g. Na is determined not only by the element content of the bedrock or other deposits from which the materials originated, but also by the effects of climatic and biological factors as well as by influences of agricultural and industrial operations that have acted on the materials for various periods of time. In the agricultural Hail soil, Na values range was 0.19-4.81 mg/g, While, the Al concentrations varied in the range of 10.26-81.26 mg/g (Table 5). The soil pH is the most critical factor controlling the amount of  $\text{Al}^{3+}$  available for plant uptake in the soil solution.

**Table 4** levels of elements in the soil of some cities in Saudi Arabia ( $\mu\text{g/g}$ )

City	Zn	Cu	Ni	Cr	Pb	Cd	Co	Se	Al
Al-Qassim region (AL-WABEL et al. 2017)	1645.40	ND-15.9	ND-14.4	8.10-28.10	6.5-149	ND-5.40	ND-3.80	-	-
Al-Hassa Oasis (MOHAMMED et al., 2014)	15.45-40.16	26.61-57.33	4.32-10.14		2.77-4.14	0.04-0.08	2.46-4.14		
Northwestern, Saudi Arabia (NAZZAL et al. 2016)	10.9-52.2	5.8-42.8		14.8-37.6	15.9-62.4	0.09-0.78	0.9-4.9		
Al-Kharj region, (AL-HAMMAD & ABDEL-SALAM, 2016)	38.45- 174.52	-	14.70- 49.52	43.50- 89.23	18.71- 42.85	0.194- 0.475	21.87-91.34	-	-
Gulf of Aqaba (GHREFAT et al. 2016)	-	-	-	-	-	-	-	0.1- 3	0.5-6.8
Al-Hayr area- Riyadh (HASAYAN et al. 2017)	9.52-27.40	8.86-10.91	11.13-19.23	10.48-20.30	5.60-7.14	0.06-0.16	-	-	-
Present study	43.29-1293.31	8.44-105.51	3.67-86.93	6.28-140.05	1.66-90.31	0.12-2.09	0.86-48.27	0.40-8.90	0.72-245.6

**Table 5** Summary of major elements concentration ( $\mu\text{g/g}$ ) in Hail soil

Variable	Na	Mg	Ca	Al
Max	117.19	74.16	39.35	245.57
Min	1.90	2.34	0.19	0.72
Median	9.54	7.03	0.69	18.16
Mean	15.12	9.40	2.54	27.28
SD	18.70	10.77	5.97	35.90

Calcium and Magnesium are secondary nutrient and plants require them in quantities as phosphorus. Where, Ca and Mg ions held to the surface of clay and organic matter in the soil by electrostatic charge. When Ca and Mg are abundant in the soluble phase tree roots absorb these nutrients by mass flow. If Ca and/or Mg are less abundant or limited by soil moisture, uptake occurs more slowly through diffusion. The Ca and Mg in soil varied considerably in Hail area, where the ranges were 1.90-37.99 and 2.34-74.16mg/g, respectively (Table 5). For Na, Al, Ca and Mg, no significant differences were observed between all soil samples.

### 3.4 Contamination level of elements

#### 3.4.1 *I-geo* values

The result of *I-geo* values in the soil samples were presented in Table 6 and were mostly fell on the negative site, where, the ranges were not very wide. *I-geo* values for Zn showed that 43% of the samples fell in the unpolluted class, 28% in the unpolluted–moderately polluted category, 26% are somewhat polluted and 4% are moderated to strongly polluted. *I-geo* values for Cu, Ni and Cr showed that more than 89% of the samples fell in the uncontaminated class. While *I-geo* for Pb, Cd, and Co showed that 100% of the samples fell in the unpolluted category. Finally, *I-geo* for Se showed that 2% of the samples dropped in the uncontaminated class and 24% in the unpolluted-moderately polluted level, 46% are moderately polluted, and 28% are moderated to strongly polluted. *I-geo* values in the analyzed soils revealed no real sign of contamination with almost all the samples reflecting a lack of contamination for all elements except Se and Zn, which showed a moderate level of pollution due to anthropogenic sources. These are suggested the absence of the variety of soil features and pollution sources in the studied area [2].

**Table 6** *I-geo* value of trace elements in Hail soil

Variable	Max	Min	Mean	Median	SD	Soil quality
Na	15.77494	9.491853	11.71383	11.98371	1.724339	Extremely polluted
AS	-3.906890596	-3.906890596	-3.906890596	-3.906890596	0	Unpolluted
Cd	-2.03686	-10.0653	-7.24821	-7.08164	2.494175	Unpolluted
Pb	7.909453	3.881906	5.910909	5.883743	.945549	Extremely polluted
Cr	7.78017	3.752623	5.781626	5.75446	0.945549	Extremely polluted
Ni	8.736791	5.07893	7.140423	7.360408	0.995346	Extremely polluted
Fe	24.03171	21.23189	23.02289	23.20777	0.716055	Extremely polluted
Mn	14.16601	10.48543	13.00226	13.46438	0.988808	Extremely polluted
Ba	10.46557	7.44427	9.467148	9.788827	0.824032	Extremely polluted
Cu	8.316037	4.597585	6.807031	6.86051	0.938145	Extremely polluted
Zn	13.2972	7.184941	11.22702	11.91099	1.802304	Extremely polluted

### 3.5 Nemerow pollution index

The values, range, mean and SD for  $P_i$ , Nemerow P and PLI in the samples were shown in Table 7. The single index ( $P_i$ ) clarified that the Zn and Se pollution intensity was strong, where the pesticides and fertilizers were the well-known external sources of agricultural soil elements in addition to natural causes [36, 37]. PLI did not show much fluctuation, where 4.2% of samples showed high levels (>1). Lower values of PLI imply no considerable input from anthropogenic sources. Nemerow P for Zn, Cu, Ni, Cr, Se and Cd was more than 1 indicating slight overall pollution, while for Co and Pb, it was less than 1, showing no contamination. The levels of elements in agricultural soils are mainly affected by parent materials in addition to pesticide and fertilizer application [36, 37].

**Table 7** Descriptive statistics of soil trace elements pollution indices

	Variable	Max	Min	Mean	SD	P
<i>pi</i>	Na	41.5333	0.53333	5.35873	9.11276	1332.536
	K	-	-	-	-	636.4845
	AS	0.0001	0.0001	0.0001	0	0.001
	Cd	0.00251		0.00026	0.00055	0.185625
	Pb	0.0736	0.00451	0.02247	0.01544	3.808918
	Cr	0.24555	0.03023	0.10312	0.05528	12.05233
	Ni	0.31601	0.02504	0.12806	0.07823	10.84969
	Fe	1.029	0.14777	0.56515	0.22927	4150.654
	Mn	0.68933	0.05376	0.36573	0.17759	110.3567
	Ba	0.2121	0.02612	0.12128	0.05658	17.27637
	Cu	0.12044	0.00915	0.05054	0.02886	5.818741
	Zn	0.37748	0.00546	0.1457	0.10696	57.22162
<b>PLI</b>		0.08593	0.01364	0.0452	0.02077	

### 3.6 Descriptive statistics of the EF values of soil trace elements

**Table 8** EF value of trace elements in Hail soil mg/kg

	Max	Min	Mean	SD	Pollution level
Na	46.6824	1.02138	8.84561	11.2305	significant
As	$6.77 \times 10^{-5}$	$9.72 \times 10^{-6}$	$2.24 \times 10^{-5}$	$1.42 \times 10^{-5}$	Clean light
Cd	0.0056	1.54219	0.000496	0.001198	Clean-light
pb	0.12157	0.0121	0.04292	0.02571	Clean-light
Cr	0.58686	0.05145	0.19782	0.1107	Clean-light
Ni	0.40552	0.06922	0.21958	0.07909	Clean-light
Fe	1	1	1	0	Clean-light



Mn	1.97499	0.32175	0.65946	0.35329	Clean-light
Ba	0.5077	0.097	0.22116	0.0913	Clean-light
Cu	0.1569	0.05064	0.08657	0.02748	Clean-light
Zn	0.68544	0.01741	0.24658	0.17271	Clean-light

Table 8 summarized the descriptive statistics of the EF values of soil trace elements. Where, the highest EF values (mean > 20) of soil trace elements recorded for Zn and Cd, which indicated substantial pollution influenced by anthropogenic sources [36,37]. The average EF value of Cu, Ni and Pb were more than 5, suggesting that they had been impacted by natural or anthropogenic sources. The average EF values of Mn, Cr, and Co were comparable; their EF values were less than 5, where, the soil sampling sites had moderate pollution. Finally, for Fe the EF values were less than two suggested that the elements come entirely from crustal materials or natural processes.

### 3.7 Aotal Dissolved Solids (TDS)

The total concentrations of soluble salts in water have great effects on plants and different crops. If the salt concentration in water increases, it is difficult for plant to extract water. The irrigation water quality classification schemes have been proposed by many researchers. Those classifications depend on one or more factors; one of the most important factors is salinity of water which expresses by TDS. Wilcox and Magistad (1943) adopted more simplified classification as given in table (8). Also, another classification is there as shown in table (9), according to (FAO, 1985). [38, 39].

**Table 9** Standards for irrigation water [38, 39]

Water class	EC (Mhos/cm <sup>-1</sup> )	Salt content (mg/l)	Sodium (%)	Boron (ppm)
Class 1	1000	700	60	0.5
Class 2	1000-3000	700-2000	60-75	0.5-2.0
Class 3	>3000	>2000	>75	>2.0

Class 1: Excellent to good, suitable for most plants most conditions; Class 2: Good to injurious, probably harmful to more sensitive crops; Class 3: Injurious to unsatisfactory, probably harmful to most crops.

## 4. Conclusion

From the present experimental study, we can have concluded the following point

- Agriculture soil contamination were studied using analytical calculation and comparison using ICP-MS and flame photometer technique.
- The mean of the elements concentrations in irrigating water were found as the following order: Na > K > Mn > Ba > Fe > Zn > Pb > Cr > Ni > Cu > Cd > As.
- The mean of the elements concentrations in soil were found as the following order: Na > K > Fe > Mn > Zn > Ba > Cr > Ni > Cu > Pb > Cd > As.
- the mean values of Cr, Ni, Cu, Pb, Cd, and As, did not exceed the Canadian soil guidelines, which mean that the soil is not contaminated. Zn were present in high levels and had a significant contamination level compared to other elements.
- Nemerow P, *I-geo*, and PLI indices were successfully applied for the assessment of elemental contamination of Hail soil. However, the highest EF values (mean > 20) of soil trace elements was recorded for Zn and Cd, which indicated heavy pollution influenced by anthropogenic sources.
- The EF values for Se were less than two suggested that the element comes mainly from a geological source; however, the anthropogenic activities related to use of fertilizers in the agricultural lands may have led to an increased amount of this element in the soil.

- Consequently, the risk of Cd and Zn accumulation in Hail soil requires further attention and monitoring to make sure the agrochemical inputs are responsible for the high accumulation of elements in soil.
- The used irrigation water as soon as agriculture soil at Hail region Saudi Arabia are suitable and considered as a good soil comparison with the international standards.

---

## References

- [1] Bhattacharya P, Misra S, Hussain M. Nutritional Aspects of Essential Trace Elements in Oral Health and Disease: An Extensive Review. *Scientifica*. 2016; 1-12.
- [2] Abdel-Satar A, Al-Khabbas M, Alahmad W, Yousef W, Alsomadi R, Iqbal T. Quality assessment of groundwater and agricultural soil in Hail region, Saudi Arabia. *The Egyptian Journal of Aquatic Research*. 2017; 43: 55-64.
- [3] Yahaya Y, Birnin-Yauri UA, Bagudo BU, Noma SS. Quantification of macro and micro elements in selected green vegetables and their soils from Aliero agricultural fields in Aliero, Kebbi State, Nigeria. *Journal of Soil Science and Environmental Management*. 2012; 3: 207-215.
- [4] Farghaly OA, Ali AMM, Ghandour MA. *Egypt J. Anal. Chem.* 1999; (8): 70- 83.
- [5] Martinotti W, Queirazza G, Guarinoni A, Mori G, *Anal. Chim. Acta.* 1995; (305): 183-191.
- [6] Farghaly OA, Ghandour MA. *Talanta* 1999; (49): 31.
- [7] Ouyang R, Zhu Z, Tatum CE, Chambers JQ, Xue ZL. *Journal of Electroanalytical Chemistry*. 2011; 656(1-2): 78–84.
- [8] Thanh NN, Hop NV, Luyen ND, Phong NH, Toan TTT. *Advances in Materials Science and Engineering*. 2019; 11.
- [9] US Environmental Protection Agency (US EPA). Supplementary Guidance for Conducting Health Risk Assessment of Chemical Mixtures. EPA/630/R-00/002. Aug 2000.
- [10] WHO (World Health Organization). Guidelines for drinking-water quality, FIRST ADDENDUM TO THIRD EDITION Volume 1 Recommendations, World Health Organization, Geneva. 2017.
- [11] He Z, Yang X, Stoffella P. Trace elements in agroecosystems and impacts on the environment. *Journal of Trace Elements in Medicine and Biology*. 2005; 19: 125-140.
- [12] Hereher M, Al-Shammari A, Abd Allah S. Land Cover Classification of Hail—Saudi Arabia Using Remote Sensing. *International Journal of Geosciences*. 2012; 03: 349-356.
- [13] Farghaly OA, Abdel Hameed RS, Abd-Alhakeem H Abu-Nawwas. Analytical Application Using Modern Electrochemical Techniques. *International Journal of Electrochemical Science*. 2014; 9: 3287 – 3318.
- [14] OA Farghaly, RS Abdel Hameed, Abd-Alhakeem H Abu-Nawwas. Electrochemical analysis techniques: a review on recent pharmaceutical applications. *International Journal of Pharmaceutical Science Review and Research*. 2014; 25(2): 37-45.
- [15] Reda S Abdel Hameed, Meshari M Aljohani, Ayham Bani Essa, Azaa Khaled, Amr M Nassar, Magd M Badr, Saedah R Al-Mhyawi, Mahmoud S Soliman. Electrochemical Techniques for Evaluation of Expired Megavit Drugs as Corrosion Inhibitor for Steel in Hydrochloric Acid, *Int. J. Electrochem. Sci.* 2021; 16: 1-13.
- [16] Reda S Abdel Hameed, MT QURESHI, AM AL-BONAYAN, SR AL-MHYAWI, MFH ABD EL-KADER. CORROSION INHIBITION OF STEEL IN MARINE ENVIRONMENT USING AMINO AMIDE DERIVED FROM PET PLASTIC WASTE, *Journal of Optoelectronic and Biomedical Materials*. 2021; 13(1): 1 – 10.
- [17] Reda S Abdel Hameed, AM Al-bonayan. Recycling of Some Water Soluble Drugs for Corrosion Inhibition of Steel Materials: Analytical and Electrochemical Measurements, *Journal of Optoelectronic and Biomedical Materials*. 2021; 13(2): 45 – 55.
- [18] Reda Abdel Hameed, M Al Elaimi, MT Qureshi, A Nassar, MFH Abd el-kader, Meshari Aljohani, Enas Ismail Arafa. Green Recycling of Poly (ethylene terephthalate) Waste as Corrosion Inhibitor for Steel in Marine Environment, *Egyptian Journal of Chemistry*. 2021; 64(2): 773 – 780.
- [19] Reda S. Abdel Hameed, Muhammad Tauseef Qureshi, M Abdallah. Application of Solid Waste for Corrosion Inhibition of Steel in Different Media- A Review, *Int. J. Corros. Scale Inhib.* 2021; 10(6): 68-79.

- [20] Reda. S. Abdel Hameed, Enas H Aljuhani, Rasha Felaly, Alaa M. Munshi. Effect of Expired Paracetamol-Zn+2 System and its Synergistic Effect Towards Iron Dissolution inhibition and Green Inhibition Performance, *Journal of Adhesion science and Technology*. 2020; 36: 27-43.
- [21] RS Abdel Hameed, EH Aljuhani, AH Al-Bagawi, AH Shamroukh, M Abdallah. Study of sulfanyl pyridazine derivatives as efficient corrosion inhibitors for carbon steel in 1.0 M HCl using analytical techniques, *Int. J. Corros. Scale Inhib.* 2020; 9(2): 623–643.
- [22] Reda. S Abdel Hameed, AH Al-Bagawi, Hassan A Shehata, Ahmed H Shamroukh, M Abdallah. Corrosion Inhibition and Adsorption Properties of Some Heterocyclic Derivatives on C- Steel Surface in HCl, *Journal of bio- tribio corrosion*. 2020; 51(6): 1-11.
- [23] Amaal M Abdel-Satar. Quality assessment of groundwater and agricultural soil in Hail region, Saudi Arabia, *Egyptian Journal of Aquatic Research*. 2017.
- [24] Zaidi F, Nazzal Y, Jafri M, Naeem M, Ahmed I. Reverse ion exchange as a major process controlling the groundwater chemistry in an arid environment: a case study from northwestern Saudi Arabia. *Environmental Monitoring and Assessment*. 2015; 187: 607.
- [25] SESD. U.S. Environmental Protection Agency-Science and Ecosystem Division, Operating Procedure for Groundwater Sampling, *SESDPROC-301-R3*. 2013.
- [26] Kramer R. The sinister attribution error: Paranoid cognition and collective distrust in organizations. *Motivation And Emotion*. 1994; 18: 199-230.
- [27] Jackwerth E, Wurfels M. Der Druckaufschluß – Apparative Möglichkeiten, Probleme und Anwendungen. In: Stoeppler, M. (Ed.), *Probennahme und Aufschluß*. Springer-Verlag, Berlin. 1994; 121–138.
- [28] Tomlinson D, Wilson J, Harris C, Jeffrey D. Problems in the assessment of heavy-metal levels in estuaries and the formation of a pollution index. *Helgoländer Meeresuntersuchungen*. 1980; 33: 566-575.
- [29] Cheng J, Shi Z, Zhu Y. Assessment and mapping of environmental quality in agricultural soils of Zhejiang Province, China. *Journal of Environmental Sciences*. 2007; 19: 50-54.
- [30] Atiemo S, Ofosu F, Aboh I, Oppon O. Levels and sources of heavy metal contamination in road dust in selected major highways of Accra, Ghana. *X-Ray Spectrometry*. 2012; 41: 105-110.
- [31] Zhang J, Yang R, Chen R, Peng Y, Wen X, Gao L. Accumulation of Heavy Metals in Tea Leaves and Potential Health Risk Assessment: A Case Study from Puan County, Guizhou Province, China. *International Journal of Environmental Research and Public Health*. 2018; 15: 133.
- [32] He Z, Yang X, Stoffella P. Trace elements in agroecosystems and impacts on the environment. *Journal of Trace Elements in Medicine and Biology*. 2005; 19: 125-140.
- [33] Hereher M, Al-Shammari A, Abd Allah S. Land Cover Classification of Hail—Saudi Arabia Using Remote Sensing. *International Journal of Geosciences*. 2012; 03: 349-356.
- [34] Pakade V, Cukrowska E, Chimuka L. Comparison of antioxidant activity of *M. oleifera* and selected vegetables in South Africa. *South African Journal of Science*. 2013; 109: 1-5.
- [35] Kebir T, Bouhadjera K. Heavy metal concentrations in agricultural soils and accumulation in plants growing near of dumpsite of Ghazaouet (west of Algeria). *International Journal of Current Research*. 2011; 2: 42–49.
- [36] Wei B, Yang L. A review of heavy metal contaminations in urban soils, urban road dusts and agricultural soils from China. *Microchemical Journal*. 2010; 94: 99-107.
- [37] Szolnoki Z, Farsang A, Puskás I. Cumulative impacts of human activities on urban garden soils: Origin and accumulation of metals. *Environmental Pollution*. 2013; 177: 106-115.
- [38] Artemio S, Ofosu F, Abloh I, Oppon O. Levels and sources of heavy metal contamination in road dust in selected major highways of Accra, Ghana. *X-Ray Spectrometry*. 2012; 41: 105-110.
- [39] Cheng J, Shi Z, Zhu Y. Assessment and mapping of environmental quality in agricultural soils of Zhejiang Province, China. *Journal of Environmental Sciences*. 2007; 19: 50-54.