

(RESEARCH ARTICLE)



The role of computed tomography in diagnosis of patients with chest trauma

Hilendarov A*, Al Georgiev and L Chervenkov

Department of Diagnostic Imaging, Medical University, Plovdiv, Bulgaria.

International Journal of Frontline Research in Life Science, 2023, 02(01), 033–042

Publication history: Received on 03 November 2023; revised on 23 December 2023; accepted on 26 December 2023

Article DOI: <https://doi.org/10.56355/ijfrls.2023.2.1.0046>

Abstract

Purpose: Chest traumas are a significant cause of mortality and morbidity. The purpose of this article is to present the place of CT in the diagnosis of chest trauma patients.

Methods and Materials: Multi detector computed tomography (MDCT) is considered to be the most effective imaging method in trauma patients. Its advantages include high speed and high geometric resolution at any plane. The method allows us to view large parts of the body with minimal motion artifacts and to create accurate multiplane reformations. One hundred and twelve patients are examined at the Emergency department. The injuries of the chest and intrathoracic organs are found. The research method is discussed by imaging diagnosticians and clinicians with a view to using the most informative diagnostic method in each individual case. Of course, informed consent is also taken from the patients. The MDCT machine are used predominantly in these patients.

Results: One hundred and twelve patients are examined at the Emergency department in purpose to evaluate chest lesions. MDCT allows us to view diagnosis significantly more accurate procedure. The injuries of the chest and intrathoracic organs are found. In 56 (50%) of all 112 patients included in the study, fractures of the ribs are the most common traumas. Also fracture of the scapula, sternum, extra pleural hematomas, pneumothorax, hemothorax, pulmonary contusions, laceration of trachea and bronchi, rupture of esophagus, diaphragmatic rupture and injuries of blood vessels are established.

Conclusion: We present our experience in imaging diagnosis and behavior in poly trauma patients in conditions of emergency. MDCT must be used as first necessary investigation a procedure which allows us to view diagnosis significantly more accurate than other imaging methods.

Keywords: Computed tomography; Thorax; Penetrating trauma; Traumatic injuries; Emergency patients

1. Introduction

In recent years has been observed major chest injuries witch represent a significant socio-economic problem. They commonly occur in the younger population and are the most common cause of death in people between 25 and 40 years of age. Chest injuries occur in about 20 % of all trauma patients. In up to 80 % of cases they are associated with injuries of other body parts such as the head (69%), abdomen and pelvis (43%) and extremities (52%) [1, 2]. Diagnostic imaging plays a key role in deciding the therapeutic procedure [3]. Multi detector computed tomography (MDCT) is considered to be the most effective imaging method in this field and therefore should be a part of the diagnostic process.

We are trying to evaluate the place of CT in the diagnostic algorithm in major trauma patients.

* Corresponding author: Hilendarov

2. Material and methods

Clinical symptoms of chest trauma often do not correlate with their severity. This is the reason why diagnostic imaging are the first procedures performed after admission to the emergency department. The simplest and fastest methods include chest radiography and ultrasound. They can provide important information about the presence of serious injuries requiring an emergency intervention, such as tension pneumothorax, large hemothorax, hemothorax and injuries of the abdominal organs. It is also very unreliable in detecting injuries of the heart and great vessels [4–8]. Compared to this, MDCT enables sufficiently accurate assessment of all compartments of the chest and also reveals changes that are not detectable by other methods (Fig. 1).




Figure 1 a, b Patient with laceration of the aorta. A/ Tomogram (analogical to chest radiograph) shows normal mediastinal width. B/ MDCT reveals rupture of the aorta with small hematoma (arrow)

Excellent spatial and temporal resolution is the main benefit. It allows performing two-dimensional (2D) and three dimensional (3D) reformations in any plane and angle of view.

The choice of a diagnostic procedure depends on the patient's condition and traumatic mechanism. In stable patients with no need of emergency intervention, the MDCT examination can be performed directly after admission to emergency department and primary clinical examination. In the case of circulatory instability the chest X-ray in supine position is usually performed as well as ultrasound examination to exclude findings that require an immediate intervention.

Generally speaking, in order to minimize motion artifacts high examination speed should be used in traumas, and because of the need to perform reformations in other planes the highest resolution in the Z-axis should be used as well. (Fig. 2) [6].




Figure 2 Low-dose examination. Significant dose reduction was achieved (80 KV, 40 effective maps, dose-length product 37 mGr *cm). Coronal edge enhancement reconstruction

Application of contrast media is essential for the assessment of vascular structures and parenchymal organs and for detection of active bleeding. Therefore, we used a longer scanning delay (30–40 s.) compared with standard chest examinations. A small amount of contrast material we used in case of suspected rupture of the esophagus. In this case water-soluble iodinated contrast medium (usually 5–10 % solution) was used. The slices are reconstructed in three principal planes (axial, coronal and sagittal). We used thinner slices (1– 1.5 mm) using high-resolution images.

3. Results and discussion:

Chest wall injuries are very common. Fractures reflect the intensity and direction of forces during trauma. However, the extent of a chest wall injury does not have to correlate with the injuries of the intrathoracic organs. no fractures. Patients according to the type of injuries are presented on Table1. This is evident especially in children and young adults with a flexible skeleton in whom severe traumas may occur despite there being no fractures.

Table 1 Patients according to the type of injuries

S.N	Type of injuries	Number Of patients	% of all
1.	Fractures of the ribs	56	50 %
2.	Fracture of the scapula	11	9.82%
3.	Sternal fractures	9	8.03%
4.	Extrapleural hematomas	45	40.17 %
5.	Pneumothorax	35	31.25%
6.	Hemothorax	39	34.82 %
7.	Pulmonary contusions	79	70.53 %
8.	Lung laceration	64	57.14 %
9.	Lung torsion	21	18.75 %
10.	Lung herniation	34	30.35 %
11.	Laceration of trachea and bronchi	59	52.67 %
12.	Rupture of esophagus	13	11.60 %
13.	Diaphragmatic rupture	7	6.25 %
14.	Large chest vessels	45	40.17 %
15.	Other blood vessels	28	25 %
16.	Heart contusion	32	28.54 %

The total number and percentages are not equal to the total number of patients because one patient has several types of injuries.

In our study, fractures of the ribs are the most common chest wall traumas. In 56 (50%) of all 112 patients included in our study, this is the most common chest wall traumas. They in fact are usually not clinically significant. MDCT determine the number of fractures, their location and degree of dislocation more accurately than X-rays. Contrary to X-rays, it is also able to detect fractures of the rib cartilages [3, 5].

In eleven (9.82%) of the cases a fracture of the scapula was established. It is usually caused by strong direct impact or the large axial forces. In up to 40 % of cases, it is associated with pulmonary contusion, pneumothorax or hemithorax [9- 11]. The areas of the scapular body and neck are mostly affected (Fig. 3).



Figure 3 Volume-rendering image. Fracture of the right scapula and multiple rib fractures on the right

Sternal fractures occur in 9 (8.03%) of cases. They are usually caused by direct impact of the anterior chest wall (typically impact with the steering wheel). Anterior dislocation is more common and clinically less severe. Posterior dislocations may be associated with major vascular traumas [12-13]. In the MDCT examination we can find blood collection separated from the lungs and pleural cavity by a thin strip of fat (Fig. 4) [14]



Figure 4 Unstable dislocated fracture (white arrows), combined with dislocation in the sternal angle (black arrow) and retrosternal hematoma




Figure 5 Combination of extra pleural hematoma and hemi thorax. Extra pleural hematoma is surrounded by a thin strip of fat (arrow)

In forty five (40.17%) of patients extra pleural hematomas are caused by injury of the intercostal artery. They occurs relatively rarely as a result of the chest wall injury or a complication of interventions (drainage, insertion of the central

venous catheter). Blood accumulates between the parietal pleura and endo thoracic fascia. In the MDCT examination we find blood collection separated from the lungs and pleural cavity by a strip of fat (Fig. 5) [14].

In 35(31.25%) of all cases a pneumothorax accompanies blunt and penetrating chest trauma. It is the second most common finding after rib fractures. It is caused by direct injury to the pleura, but also by a rupture of the alveoli accompanied by increased pressure in the airways. On CT it is manifested as a collection of gas in the pleural cavity. In chest Xray examinations performed in the supine position, it is often manifested very discretely, and it cannot be seen at all (Fig. 6) [15,16].



Figure 6 Pneumothorax: MDCT shows significant pneumothorax in the ventral part of the left pleural cavity

In 5(4.46%) of the patients tension pneumothorax is a serious, life-threatening condition requiring immediate drainage. It is often clinically diagnosed before MDCT. Its basic symptoms include an increase of the volume of the affected hemithorax, shift of the mediastinum to the healthy side, and compression of the mediastinal vessels (mainly veins), heart and depression of the diaphragmatic arch (Fig. 7).




Figure 7 Tension pneumothorax: Maximum intensity projection: mediastinal shift to the right, dislocation and compression of the heart, and compression of the left pulmonary artery

In 39(34.82%) of all cases with hemithorax, bleeding into the pleural cavity is most commonly caused by laceration of the pulmonary parenchyma and pleural injury. In 12(10.71%) of cases the injury to an artery (usually intercostal), bleeding progresses rapidly and requires surgical therapy. MDCT is the method of choice for diagnosis of pleural fluid even better than US. The density of liquid blood is between 30 and 50 HU, and the density of blood clots is 50–90 HU. Hematomas can sometimes have a layered structure (Fig. 8).



Figure 8 Large layered hemi thorax. Hyper dense clots (50 HU) in the dorsal part of the right pleural cavity (arrow).
Mediastinal shift to the left side

Injury of the lung parenchyma includes pulmonary contusion, laceration, torsion and herniation. These conditions can be complicated by atelectasis, aspiration pneumonia or acute respiratory distress syndrome (ARDS) [6]. Lung trauma is a reason for surgical treatment in 32(28.57%) of the patients. This is indicated in cases of injury of the major blood vessels, signs of active bleeding, large hematomas or hemodynamic instability

Pulmonary contusions was found in 79 (70.53%) of all 112 cases and is the most common trauma of the lung parenchyma. It can be detected on the X-ray image with up to a 6- h delay. On the other hand, CT is very sensitive to contusion changes, and it can detect them earlier and more accurately and allows their quantification, which is important in the determination of further treatment. In contusions exceeding 20–30 % of the total lung volume, mechanical ventilation is indicated to prevent the development of complications, especially acute respiratory distress syndrome [17- 19].

The CT findings depend on the severity of parenchymal affection. Simultaneously, parenchymal lacerations can occur (Fig. 9). The peripheral areas of the lung are affected, and the distribution does not respect boundaries of the lobes and segments [6].




Figure 9 Lung contusion with small lacerations (arrow)

X-ray examination is usually negative because of superposition of simultaneously occurring contusion Lung laceration occurs in 64 (57.14%) penetrating injuries as a result of direct shearing forces. . At present we can find it relatively commonly because of the broad use of CT in traumatized patients changes. If a laceration reaches the pleural cavity, it can cause pneumothorax or hemothorax. Laceration closed in the parenchyma has the appearance of a thin-walled, spherical, oval or tubular cavity containing air, blood or their combination (Fig. 10). There are usually contusion changes around it. When they are absorbed the laceration, which does not contain air, can mimic a pulmonary node [6].




Figure 10 a, b. Extensive lung laceration. a/ Cavities with air-fluid levels in the right upper lobe, right-sided pneumothorax, bilateral subcutaneous emphysema, pneumonic- mediastinum, mediastinal shift to the left. b/ Extravasation of the contrast material into the cavity due to active bleeding

Lung torsion in 34 (30.35%) patients occurs a lung herniation in a rare complication of chest injuries. It occurs at the site of a defect of the chest wall with a significant increase in intrathoracic pressure. The chest wall defects are caused by multiple fractures of the ribs and clavicular dislocation. If parenchymal tearing occurs at the same time, the herniation may be associated with pneumothorax and hemi thorax [20-23].

Rupture of the esophagus is presented in 13(11.60%) patients as a part of a blunt or penetrating chest trauma . Most often, it occurs in the cervical and upper mediastinal part. It is usually caused by bone fragments from the injury of adjacent sections of the spine. The CT findings include thickening of the esophageal wall, pneumomediastinum and para esophageal fluid collections [24, 25].

Diaphragmatic rupture occurs in 7(6.25%) of patients hospitalized with a major trauma [26]. It is a frequently overlooked injury but it is clinically very serious. Laceration in the right side is more difficult to detect. Sensitivity of the initial X-ray examination is low, 17 % and 27–60 % on the right and left side, respectively. Compared to this, MDCT reaches a sensitivity of 50–83 % and 78–100 % in the right and left side, respectively. Evaluation of multiplane reformations in the coronal and sagittal plane is very important for making the diagnosis. The CT signs of a diaphragmatic rupture include discontinuity of the diaphragm with thickening at the edge of the defect, herniation of the abdominal organs into the chest, strangulation of the organs at their passage through the diaphragmatic defect (collar sign) and location of herniated abdominal organs in the dorsal part of the thorax where they are in contact with the chest wall (dependent viscera sign) (Fig. 11). Diaphragmatic injuries are often associated with injuries of the spleen, liver, lungs and rib fractures.



Figure 11 Herniation of abdominal organs into the left hemi thorax (arrow), club shaped edge of the ruptured diaphragm (arrowhead)

Trauma of the large chest vessels represents in about 45 (40.17%) of all major trauma patients, but its incidence is probably underestimated. Although relatively infrequent, it is the second most common cause of death after head injuries. The aorta is the most commonly affected vessel. Rarely, we can see injuries of the aortic branches, internal thoracic artery or major mediastinal veins. In the case of a blunt trauma, there are three traumatic mechanisms leading to the trauma—deceleration, torsion and increased intraluminal pressure. Findings on the X-ray examination are usually nonspecific and do not allow to differentiate trauma of the large vessels from other less serious conditions with similar X-ray findings, as described above. [5]. If this type of trauma is suspected, MDCT is currently the first-choice method. Its sensitivity and specificity are almost 100 %. Angiography is indicated only in cases of doubt and for navigation of the interventional procedures.

Mortality of untreated aortic trauma exceeds 95 %. Those who come to the hospital have a 70 % chance of survival if the trauma is quickly diagnosed and treated. Treatment is either surgical or radio interventional (endovascular stent grafting). The most common site of injury (approximately 90 %) is the isthmus, which represents the transition between the relatively mobile arch and fixed descending part. A complete wall rupture (transection) usually results in immediate death. Direct signs of aortic trauma are pseudo aneurysm (Fig. 12), direct extravasation of contrast medium into the mediastinum, intimal flap, irregular contour or mural thrombus.




Figure 12 Aortic isthmus laceration. Pseudo aneurysm in a typical location (arrow), mediastinal hematoma, bilateral hemothorax

In 28(25%)of patients injuries of other blood vessels, such as the internal thoracic artery, branches of the aortic arch, pulmonary artery or mediastinal veins, are less common than aortic trauma, but their severity is similar. In case of bleeding from the internal thoracic artery, a hematoma occurs in the anterior mediastinum, and it may compress the right ventricle. Injury of the branches of the aortic arch occurs most frequently in the traction mechanism in case of abnormal extension of the neck or pulling of the arm.

Rare injuries of the venous structures are manifested as hematoma with extravasation of contrast medium (Fig. 13). The mechanism is similar to injuries of the arteries. Injury of the vena cava superior can spread to the right atrium as well.




Figure 13 Rupture of the azygos vein with contrast material leakage into the large hemothorax (white arrow) combined with aortic pseudo aneurysm (black arrow)

Contusion is the most common heart injury in 32(28.54%) of our cases affecting preferably the right ventricle. It is generally radiologically silent and causes arrhythmia or other electrocardiographic changes. Because of its very high mortality, cardiac rupture is rarely diagnosed by imaging methods. Diagnosis of hemo pericardium using X-ray examination is extremely problematic. The finding can be negative even in cases of cardiac tamponade. In contrast, the CT examination as well as ultrasound is a highly sensitive method. Cardiac tamponade is manifested by flattening and a reduction of the cardiac parts (Fig. 14). More rarely, we can see pneumonic pericardia, which, if large, may also result in cardiac tamponade. Herniation of the heart is a serious complication of traumatic rupture of the pericardium.




Figure 14 Hem pericardium with heart tamponade: compression of all heart parts

4. Conclusion

Our experience and literature report suggest that imaging methods are an mandatory part of the diagnostic algorithm in major chest traumas. MDCT is the main examination. It shows traumatic changes quickly, accurately and clearly allows their classification. Therefore a traumatic radiologist becomes a member of the team making decisions about the therapeutic process.

Compliance with ethical standards

Disclosure of conflict of interest

No conflict of interest to be disclosed.

Statement of ethical approval

The performed diagnostic procedures was approved by Ethical Board of Medical University

Statement of informed consent

Informed consent was obtained from all participants included in the study, before any diagnostic procedures. That was approved by Ethical board of Medical University

References

- [1] Watermark M, Schneider P (2004) MDCT of chest trauma. In: Schlep UJ 9 (ed) Multi detector-row CT of the thorax. Springer, Berlin, Heidelberg, New York, pp 409–421
- [2] Loci cero J, Mattox KL (1989) Epidemiology of chest trauma. Surge Clin North Am 69:15–19
- [3] Lomo schists FM, Eisenhower E, Lindau KF, Palo Scheck P, Schroder M, Balkier AA (2003) Imaging of chest trauma: radiological patterns of injury and diagnostic algorithms. Euro J Radio 48:61–70
- [4] Scaglione M, Pinto A, Pedraza I, Sparano A, Romano L (2008) Multi-detector row computed tomography and blunt chest trauma. European J Radio 65:377–388

- [5] Kaela R, Avery LL, Arsan AV, Novel line RA (2008) Multi detector CT of blunt thoracic trauma. *Radio Graphics* 28:1555–1570
- [6] Emalia M, Badin A, Mural MS, Arsan B, Ceyhan M, Germen N (2007) Lung parenchymal injury and its frequency in blunt thoracic trauma: the diagnostic value of chest radiography and thoracic CT. *Deign Interv Radio* 13:179–182
- [7] Brink M, Deink J, Dekker HM et al (2008) Added value of routine chest MDCT after blunt trauma: evaluation of additional findings and impact on patient management. *Am J Roentgen* 190:1591–1598
- [8] Brink M, Kool DR, Dekker HM et al (2009) Predictors of abnormal chest CT after blunt trauma: a critical appraisal of the literature. *Clink Radio* 64:272–283 traumas. *Pediatric Radio* 39:485–496
- [9] Tack D (2010) Radiation dose optimization in thoracic imaging. *JBR-BTR* 93:15–19
- [10] Kappamaki VV, Kuru MJ, Koskinen SK (2004) Multi detector CT in shoulder fractures. *Emerge Radio* 11:89–94
- [11] McGinnis M, Denton JR (1989) Fractures of the scapula: a retrospective study of 40 fractured scapulae. *J Trauma* 29:1488–1493
- [12] Kim EY, Yang HJ, Sung YM, Hwang KH, Kim JH, Kim HS (2011) Sternal fracture in the emergency department: diagnostic value of multi detector CT with sagittal and coronal reconstruction images. *Euro J Radio* 14 [Epic ahead of print]
- [13] Maier D, Jaeger M, Izadpanah K, Bornebusch L, Südkamp NP (2011) Traumatic injuries of the sternoclavicular joint. *Unfallchirurg* 114:611–621
- [14] Aquino SL, Chiles C, Oaks T (1997) displaced extra pleural fat as revealed by CT scanning: evidence of extra pleural hematoma. *Am J Roentgen* 169:687–689
- [15] Ball CG, Kirkpatrick AW, Lapland KB et al (2005) Incidence, risk factors, and outcomes for occult pneumothoraxes in victims of major trauma. *J Trauma* 59:917–924
- [16] Omar HR, Abdelmalak H, Manger D, Rashad R, Hellas E, Camporees EM (2010) Occult pneumothorax, revisited. *J Trauma Manage Outcome* 4:3–6
- [17] de Moya MA, Manolakaki D, Chang Y et al (2011) Blunt pulmonary contusion: admission computed tomography scan predicts mechanical ventilation. *J Trauma* 71:1543–1547
- [18] Strumwasser A, Chu E, Yeung L, Microflora E, Sajida J, Victorian GP (2011) A novel CT volume index score correlates with outcomes in polytrauma patients with pulmonary contusion. *J Surg Res* 170:280–285
- [19] Wong PS, Goldstar P (1992) Pulmonary torsion: a questionnaire survey and a survey of the literature. *Ann Thoracic Surge* 54:286–288
- [20] Gilkison RC, Lange P, Kirby TJ (2000) Lung torsion after lung transplantation evaluation with helical CT. *Am J Roentgen* 174:1341–1343
- [21] Sukkarieh F, Vanmeerhaeghe A, Brasseur P (2002) Post traumatic intercostal lung herniation: a case report. *J Radio* 83:1085–1087
- [22] Le Guan M, Beige man C, Bouhemad B, Weidie Z, Marmion F, and Robby JJ (2007) Chest computed tomography with multiplane reformatted images for diagnosing traumatic bronchial rupture: a case report. *Crit Care* 11:R94
- [23] Rezende-Neto JB, Hoffmann J, Al Mahroos M et al (2010) Occult pneumomediastinum in blunt chest trauma: clinical significance. *Injury* 41:40–43
- [24] Trades T, Wafaizadeh A, Steinhausen E, Krakamp B, Bouillon B (2009) Extension injury of the thoracic spine with rupture of the esophagus and successful conservative therapy of concomitant mediastinitis. *Euro Spine J* 18:240–244
- [25] Young CA, Manias CO, Ballan S, Prasad SR (2008) CT features of esophageal emergencies. *Radio Graphics* 28:1541–1553
- [26] Meyers BF, McCabe CJ (1993) Traumatic diaphragmatic hernia: occult marker of serious injury. *Ann Surge* 218:783–790